




Meningococcal vaccination for children and adolescents without risk factors



Meningococcal disease is a rare but serious disease that can cause significant illness, disability and death. Meningococcal vaccination for certain groups of people is funded under the National Immunisation Program and by states and territories.

No single vaccine protects against all serogroups



Recommended vaccines

The number of doses depends on the vaccine brand and the person's age when they start the vaccination course.

	6 weeks – 23 months	2–4 years	5–14 years	15–19 years
Aboriginal and Torres Strait Islander people	<ul style="list-style-type: none"> ▶ MenB <input checked="" type="checkbox"/> ▶ MenACWY (single dose at age 12 months <input checked="" type="checkbox"/>) 	<ul style="list-style-type: none"> ▶ MenB ▶ MenACWY 	<ul style="list-style-type: none"> ▶ MenB ▶ MenACWY 	<ul style="list-style-type: none"> ▶ MenB ▶ MenACWY <input checked="" type="checkbox"/>
Non-Indigenous people	<ul style="list-style-type: none"> ▶ MenB ▶ MenACWY (single dose at age 12 months <input checked="" type="checkbox"/>) 			<ul style="list-style-type: none"> ▶ MenB ▶ MenACWY <input checked="" type="checkbox"/>

See the Australian Immunisation Handbook for more details

Healthy children and adolescents do not need booster doses unless they are in certain special risk groups.

= vaccine funded under the National Immunisation Program

See the Australian Immunisation Handbook for more details.